



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

NOTE.—Thomas S. Clarkson remarks that the solution of 321 does not satisfy the conditions of the problem, since the whole number of shares is not to exceed 200. He contends that the problem is impossible as stated, and suggests that it may have been intended that each man's share is not to exceed 200. We think the problem is stated as intended and is therefore impossible.

ED. F.

325. Proposed by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

I have a chronometer whose rate is uniform. When it indicates t_1 time at Washington I find that it is h_1 hours slow. I take it to Philadelphia and when it indicates t_2 time, the local time of Philadelphia is h_2 hours faster. I bring my chronometer back to Washington and find that when it indicates t_3 time it is h_3 hours slow. If $t_1 = 5$ A. M., $t_2 = 7$ hours, 54 minutes, $t_3 = 11$ hours, 46 minutes A. M., $h_1 = 1$ hour, $h_2 = 1 \frac{203}{900}$ hours, $h_3 = 1 \frac{7}{30}$ hours, find the difference of longitude between Washington and Philadelphia.

Solution by J. EDWARD SANDERS, Weather Bureau, Chicago, Ill. and the PROPOSER.

$$(h_3 - h_1) / (t_3 - t_1) = \text{error for one hour.}$$

$$(t_2 - t_1) (h_3 - h_1) / (t_3 - t_1) = \text{error at time } t_2 \text{ in Philadelphia.}$$

$$\therefore h_1 + t_2 + (t_2 - t_1) (h_3 - h_1) / (t_3 - t_1) = \text{time at Washington.}$$

$$h_2 + t_2 = \text{time at Philadelphia.}$$

$$h_2 + t_2 - h_1 - t_2 - (t_2 - t_1) (h_3 - h_1) / (t_3 - t_1) = \text{difference of time between the two cities} = T.$$

$$\therefore T = \frac{h_2(t_3 - t_1) + h_3(t_1 - t_2) + h_1(t_2 - t_3)}{t_3 - t_1}.$$

$$15T = L = \text{difference in longitude.}$$

$$\therefore L = \frac{15(h_1 t_2 + h_2 t_3 + h_3 t_1 - h_1 t_3 - h_2 t_1 - h_3 t_2)}{t_3 - t_1}.$$

Putting $h_1 = 1$ hour, $h_2 = 1 \frac{203}{900}$ hours, $h_3 = 1 \frac{7}{30}$ hours, $t_1 = 5$ hours, $t_2 = 7$ hours 54 minutes, $t_3 = 11$ hours 46 minutes, we get

$$L = 1 \frac{13}{60}^\circ = 1^\circ 53'.$$

Also solved by T. S. Clarkson.

326. Proposed by R. D. CARMICHAEL, Princeton University.

Is the series, of which the n th term is $\frac{1.3.5.7 \dots (2n-1)}{(n+1)! 2^n (2n+3)}$ convergent?

If so, find its sum.